

JEE Main January 2026
Question Paper With Text Solution
22 January | Shift-2

MATHEMATICS



JEE Main & Advanced | XI-XII Foundation | VI-X Pre-Foundation

Office : Piprali Road, Sikar (Raj.) | Ph. 01572-241911

Website : www.matrixedu.in ; Email : smd@matrixacademy.co.in

**JEE MAIN JANUARY 2026 | 22TH JANUARY SHIFT-2****SECTION – A**

Question ID : 860654987

1. Let the locus of the mid-point of the chord through the origin O of the parabola $y^2 = 4x$ be the curve S. Let P be any point on S. Then the locus of the point, which internally divides OP in the ratio 3: 1, is :

(1) $2y^2 = 3x$ (2) $2x^2 = 3y$ (3) $3y^2 = 2x$ (4) $3x^2 = 2y$

Ans. Official answer NTA (1)

Sol.

Question ID : 860654988

2. Among the statements

(S1) : If A(5, -1) and B(-2, 3) are two vertices of a triangle, whose orthocentre is (0, 0), then its third vertex is (-4, -7)

and

(S2) : If positive numbers 2a, b, c are three consecutive terms of an A.P., then the lines $ax + by + c = 0$ are concurrent at (2, -2), :

- (1) both are correct (2) only (S1) is correct (3) both are incorrect (4) only (S2) is correct

Ans. Official answer NTA (1)

Sol.

Question ID : 860654991

3. Let $[\cdot]$ denote the greatest integer function, and let $f(x) = f(x) = \min\{\sqrt{2}x, x^2\}$. Let $S = \{x \in (-2, 2) : \text{the function } g(x) = |x|[\cdot x^2] \text{ is discontinuous at } x\}$. Then $\sum_{x \in S} f(x)$ equals :

(1) $1 - \sqrt{2}$ (2) $2 - \sqrt{2}$ (3) $2\sqrt{6} - 3\sqrt{2}$ (4) $\sqrt{6} - 2\sqrt{2}$

Ans. Official answer NTA (1)

Sol.



Question ID : 860654986

4. Let $P(10, 2\sqrt{15})$ be a point on the hyperbola $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$, whose foci are S and S'. If the length of its latus rectum is 8, then the square of the area of $\Delta PSS'$ is equal to :
- (1) 4200 (2) 2700 (3) 1462 (4) 900

Ans. Official answer NTA (2)**Sol.**

Question ID : 860654978

5. Let α, β be the roots of the quadratic equation $12x^2 - 20x + 3\lambda = 0, \lambda \in \mathbb{Z}$. If $\frac{1}{2} \leq |\beta - \alpha| \leq \frac{3}{2}$, then the sum of all possible values of λ is :
- (1) 6 (2) 3 (3) 1 (4) 4

Ans. Official answer NTA (2)**Sol.**

Question ID : 860654981

6. If $X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$ is a solution of the system of equations $AX = B$, where $\text{adj } A = \begin{bmatrix} 4 & 2 & 2 \\ -5 & 0 & 5 \\ 1 & -2 & 3 \end{bmatrix}$ and $B = \begin{bmatrix} 4 \\ 0 \\ 2 \end{bmatrix}$, then $|x + y + z|$ is equal to :
- (1) 1 (2) 2 (3) 3 (4) $\frac{3}{2}$

Ans. Official answer NTA (2)**Sol.**

Question ID : 860654995

7. If $y = y(x)$ satisfies the differential equation $16(\sqrt{x+9\sqrt{x}})(4+\sqrt{9+\sqrt{x}})\cos y \, dy = (1+2\sin y)dx, x > 0$ and $y(256) = \frac{\pi}{2}, y(49) = \alpha$, then :

MATRIX JEE ACADEMY**Office : Piprali Road, Sikar (Raj.) | Ph. 01572-241911****Website : www.matrixedu.in ; Email : smd@matrixacademy.co.in**



(1) $\sqrt{2} - 1$

(2) $2(\sqrt{2} - 1)$

(3) $3(\sqrt{2} - 1)$

(4) $2\sqrt{2} - 1$

Ans. Official answer NTA (4)**Sol.**

Question ID : 860654982

8. Let n be the number obtained on rolling a fair die. If the probability that the system

$$x - ny + z = 6$$

$$x + (n - 2)y + (n + 1)z = 8$$

$$(n - 1)y + z = 1$$

has a unique solution is $\frac{k}{6}$, then the sum of k and all possible values of n is :

(1) 21

(2) 20

(3) 22

(4) 24

Ans. Official answer NTA (3)**Sol.**

Question ID : 860654984

9. Let C_r denote the coefficient of x^r in the binomial expansion of $(1 + x)^n$, $n \in \mathbb{N}$, $0 \leq r \leq n$. If

$$P_n = C_0 - C_1 + \frac{2^2}{3}C_2 - \frac{2^3}{4}C_3 + \dots + \frac{(-2)^n}{n+1}C_n$$
, then the value of $\sum_{n=1}^{25} \frac{1}{P_{2n}}$ equals. :

(1) 675

(2) 650

(3) 580

(4) 525

Ans. Official answer NTA (1)**Sol.**

Question ID : 860654985

10. Let S and S' be the foci of the ellipse $\frac{x^2}{25} + \frac{y^2}{9} = 1$ and $P(\alpha, \beta)$ be a point on the ellipse in the first quadrant. If

$$(SP)^2 + (S'P)^2 - SP \cdot S'P = 37$$
, then $\alpha^2 + \beta^2$ is equal to :

(1) 11

(2) 13

(3) 15

(4) 17

Ans. Official answer NTA (2)**Sol.**



Question ID : 860654979

11. Let the domain of the function $f(x) = \log_3 \log_5 \left(7 - \log_2 (x^2 - 10x + 85) \right) + \sin^{-1} \left(\frac{3x - 7}{17 - x} \right)$ be $(\alpha, \beta]$. Then

 $\alpha + \beta$ is equal to :

- (1) 10 (2) 12 (3) 9 (4) 8

Ans. Official answer NTA (3)**Sol.**

Question ID : 860654993

12. Let $f(x) = [x]^2 - [x + 3] - 3$, $x \in \mathbb{R}$, where $[\cdot]$ is the greatest integer function. Then :

- (1) $f(x) = 0$ for finitely many values of x (2) $f(x) > 0$ only for $x \in [4, \infty)$
 (3) $f(x) < 0$ only for $x \in [-1, 3)$ (4) $\int_0^2 f(x) dx = -6$

Ans. Official answer NTA (3)**Sol.**

Question ID : 860654977

13. Let f and g be functions satisfying $f(x + y) = f(x)g(y)$, $f(1) = 7$ and $g(x + y) = g(xy)$, $g(1) = 1$, for all $x, y \in \mathbb{N}$.

If $\sum_{x=1}^n \left(\frac{f(x)}{g(x)} \right) = 19607$, then n is equal to :

- (1) 6 (2) 4 (3) 7 (4) 5

Ans. Official answer NTA (4)**Sol.**

Question ID : 860654992

14. If $\lim_{x \rightarrow 0} \frac{e^{(a-1)x} + 2 \cos bx + (c-2)e^{-x}}{x \cos x - \log_e(1+x)} = 2$, then $a^2 + b^2 + c^2$ is equal to :

- (1) 7 (2) 3 (3) 5 (4) 9



Ans. Official answer NTA (1)

Sol.

Question ID : 860654976

15. The number of elements in the relation $R = \{(x, y) : 4x^2 + y^2 < 52, x, y \in \mathbf{Z}\}$ is :

- (1) 77 (2) 89 (3) 86 (4) 67

Ans. Official answer NTA (1)

Sol.

Question ID : 860654980

16. Let $S = \{z \in \mathbf{C} : 4z^2 + \bar{z} = 0\}$. Then $\sum_{z \in S} |z|^2$ is equal to :

- (1) $\frac{5}{64}$ (2) $\frac{3}{16}$ (3) $\frac{7}{64}$ (4) $\frac{1}{16}$

Ans. Official answer NTA (2)

Sol.

Question ID : 860654989

17. Let $\vec{a} = 2\hat{i} - \hat{j} + \hat{k}$ and $\vec{b} = \lambda\hat{j} + 2\hat{k}$, $\lambda \in \mathbf{Z}$ be two vectors. Let $\vec{c} = \vec{a} \times \vec{b}$ and \vec{d} be a vector of magnitude 2 in yz-plane. If $|\vec{c}| = \sqrt{53}$, then the maximum possible value of $(\vec{c} \cdot \vec{d})^2$ is equal to :

- (1) 104 (2) 208 (3) 26 (4) 52

Ans. Official answer NTA (2)

Sol.

Question ID : 860654990

18. Let L be the line $\frac{x+1}{2} = \frac{y+1}{3} = \frac{z+3}{6}$ and let S be the set of all points (a, b, c) on L, whose distance from the

line $\frac{x+1}{2} = \frac{y+1}{3} = \frac{z-9}{0}$ along the line L is 7. Then $\sum_{(a,b,c) \in S} (a+b+c)$ is equal to :

- (1) 34 (2) 6 (3) 28 (4) 40

MATRIX JEE ACADEMY

Office : Piprali Road, Sikar (Raj.) | Ph. 01572-241911

Website : www.matrixedu.in ; Email : smd@matrixacademy.co.in



Ans. Official answer NTA (1)

Sol.

Question ID : 860654983

19. If the mean deviation about the median of the numbers $k, 2k, 3k, \dots, 1000k$ is 500, then k^2 is equal to :

- (1) 4 (2) 1 (3) 9 (4) 16

Ans. Official answer NTA (1)

Sol.

Question ID : 860654994

20. The area of the region $A = \{(x, y) : 4x^2 + y^2 \leq 8 \text{ and } y^2 \leq 4x\}$ is :

- (1) $\frac{\pi}{2} + \frac{1}{3}$ (2) $\frac{\pi}{2} + 2$ (3) $\pi + 4$ (4) $\pi + \frac{2}{3}$

Ans. Official answer NTA (4)

Sol.

SECTION - B

Question ID : 860654999

21. Let a vector $\vec{a} = \sqrt{2}\hat{i} - \hat{j} + \lambda\hat{k}, \lambda > 0$, make an obtuse angle with the vector $\vec{b} = -\lambda^2\hat{i} + 4\sqrt{2}\hat{j} + 4\sqrt{2}\hat{k}$ and an angle $\theta, \frac{\pi}{6} < \theta < \frac{\pi}{2}$, with the positive z-axis. If the set of all possible values of λ is $(\alpha, \beta) - \{\lambda\}$, then $\alpha + \beta + \gamma$ is equal to _____.

Ans. Official answer NTA (5)

Sol.

Question ID : 860654997

22. Let S be the set of the first 11 natural numbers. Then the number of elements in $A = \{B \subseteq S : n(B) \geq 2 \text{ and the product of all elements of } B \text{ is even}\}$ is _____.

Ans. Official answer NTA (1979)

Sol.

MATRIX JEE ACADEMY

Office : Piprali Road, Sikar (Raj.) | Ph. 01572-241911

Website : www.matrixedu.in ; Email : smd@matrixacademy.co.in



Question ID : 860654996

23. Suppose a, b, c are in A.P. and $a^2, 2b^2, c^2$ are in G.P. If $a < b < c$ and $a + b + c = 1$, then $9(a^2 + b^2 + c^2)$ is equal to _____.

Ans. Official answer NTA (9)

Sol.

Question ID : 8606541000

24. Let $[.]$ be the greatest integer function. If $\alpha = \int_0^{64} (x^{1/3} - [x^{1/3}]) dx$, then $\frac{1}{\pi} \int_0^{\alpha\pi} \left(\frac{\sin^2 \theta}{\sin^6 \theta + \cos^6 \theta} \right) d\theta$ is equal to _____.

Ans. Official answer NTA (36)

Sol.

Question ID : 860654998

25. Let $\cos(\alpha + \beta) = -\frac{1}{10}$ and $\sin(\alpha - \beta) = \frac{3}{8}$, where $0 < \alpha < \frac{\pi}{3}$ and $0 < \beta < \frac{\pi}{4}$.

If $\tan 2\alpha = \frac{3(1-r\sqrt{5})}{\sqrt{11}(s+\sqrt{5})}$, $r, s \in \mathbb{N}$, then $r + s$ is equal to _____.

Ans. Official answer NTA (20)

Sol.