

JEE Main January 2026
Question Paper With Text Solution
24 January | Shift-2

CHEMISTRY



JEE Main & Advanced | XI-XII Foundation | VI-X Pre-Foundation

Office : Piprali Road, Sikar (Raj.) | Ph. 01572-241911
Website : www.matrixedu.in ; Email : smd@matrixacademy.co.in

JEE MAIN JANUARY 2026 | 24 JANUARY SHIFT-2
SECTION - A

Question ID : 444792655

51. At 298 K, the mole percentage of $N_2(g)$ in air is 80%. Water is in equilibrium with air at a pressure of 10 atm. What is the mole fraction of $N_2(g)$ in water at 298 K?

(K_H for N_2 is 6.5×10^7 mmHg)

क

- (1) 1.23×10^{-7} (2) 1.17×10^{-4} (3) 9.35×10^{-5} (4) 9.35×10^5

Ans. Official answer NTA (3)

Sol.

Question ID : 444792661

52. The wavelength of light absorbed for the following complexes are in the order



- (I) (II) (III) (IV) (V)

क

- (1) III < IV < I < II < V (2) III < I < II < IV < V
 (3) III < I < IV < V < II (4) III < I < IV < II < V

Ans. Official answer NTA (4)

Sol.

Question ID : 444792654

53. The heat of atomisation of methane and ethane are 'x' kJmol^{-1} and 'y' kJmol^{-1} respectively. The longest wavelength (λ) of light capable of breaking the C-C bond can be expressed in SI unit as:

क

- (1) $\frac{N_A hc}{250(4y - 6x)}$ (2) $\frac{N_A hc}{250(y - 6x)}$ (3) $\frac{hc}{1000} \left(\frac{y - 6x}{4} \right)^{-1}$ (4) $N_A hc \left(y - \frac{6x}{4} \right)^{-1}$

Ans. Official answer NTA (4)

Sol.

Question ID : 444792670

54. In the Group analysis of cations, Ba^{2+} & Ca^{2+} are precipitated respectively as“

क

- (1) hydroxide & carbonate (2) carbonate & carbonate
 (3) chromate & sulphide (4) sulphide & sulphide

Ans. Official answer NTA (2)

Sol. Question ID : 444792669

55. The number of possible tripeptides formed involving alanine (ala), glycine (gly) and valine (val), where no amino acid has been used more than once is:

क


- (1) 6 (2) 4 (3) 8 (4) 3

Ans. Official answer NTA (1)


Sol.

Question ID : 444792665

56. The unsaturated ether on acidic hydrolysis produces carbonyl compounds as shown below:



Based on this, predict the solution/reagent that will help to distinguish "P" and "Q" obtained in the following reaction:



क

- (1) Fehling solution (2) Lucas reagent
 (3) 2, 4 - DNP reagent (4) Saturated NaHSO₃ solution

Ans. Official answer NTA (1)

Sol. Question ID : 444792652

57. The wavelength of spectral line obtained in the spectrum of Li^{2+} ion, when the transition takes place between two levels whose sum is 4 and difference is 2, is

क

- (1) 2.28×10^{-6} cm (2) 1.14×10^{-7} cm (3) 2.28×10^{-7} cm (4) 1.14×10^{-6} cm

Ans. Official answer NTA (4)

Sol.

Question ID : 444792658

58. The correct order of C, N, O and F in terms of second ionisation potential is

क

- (1)
- $F < N < C < O$
- (2)
- $C < N < F < O$
- (3)
- $C < F < N < O$
- (4)
- $C < O < N < F$

Ans. Official answer NTA (2)**Sol.**

Question ID : 444792668

59. Given below are two statements:

Statement I: The dipole moment of R-CN is greater than R-NC and R-NC can undergo hydrolysis under

acidic medium to produce $R - \overset{\text{O}}{\parallel}{\text{C}} - \text{OH}$ Statement II: R-CN hydrolyses under acidic medium to produce a compound which on treatment with SOCl_2 , followed by the addition of NH_3 gives another compound (x). This compound (x) on treatment with $\text{NaOCl} / \text{NaOH}$ gives a product, that on treatment with $\text{CHCl}_3 / \text{KOH} / \Delta$ produces R-NC

In the light of the above statements, choose the correct answer from the options given below

क

- (1) Both Statement I and Statement II are true
-
- (2) Both Statement I and Statement II are false
-
- (3) Statement I is false but Statement II is true
-
- (4) Statement I is true but Statement II is false

Ans. Official answer NTA (3)**Sol.**

Question ID : 444792659

60. Choose the Incorrect statement

- (1) Carbon exhibits negative oxidation states along with +4 and +2.
-
- (2)
- CO_2
- is the most acidic oxide among the dioxides of group of 14 elements.
-
- (3) Carbon cannot exceed its covalency more than four.
-
- (4) Among the isotopes of carbon,
- ^{13}C
- is a radioactive isotope.

क

Ans. Official answer NTA (4)

Sol.

Question ID : 444792660

61. "X" is an oxoanion of the lightest element of group 7 (in the periodic table). The metal is in +6 oxidation state in "X". The color of the potassium salt of X is


- (1) green (2) yellow (3) purple (4) orange

क

Ans. Official answer NTA (1)**Sol.**

Question ID : 444792657

62. Consider the following gaseous equilibrium in a closed container of volume 'V' at $^{13}CT(K)$.



2 moles each of $P_2(g)$, $Q_2(g)$ and $PQ(g)$ are present at equilibrium. Now one mole each of ' P_2 ' and ' Q_2 ' are added to the equilibrium keeping the temperature at $T(K)$. The number of moles of P_2 , Q_2 and PQ at the new equilibrium, respectively, are

क

- (1) 2.67, 2.67, 2.67 (2) 1.21, 2.24, 1.56 (3) 2.56, 1.62, 2.24 (4) 1.66, 1.66, 1.66

Ans. Official answer NTA (1)**Sol.**

Question ID : 444792664

63. Given below are two statements:

Statement I : Cross aldol condensation between two different aldehydes will always produce four different products.

Statement II : When semicarbazide reacts with a mixture of benzaldehyde and acetophenone under optimum pH, it forms a condensation product with acetophenone only.

In the light of the above statements, choose the correct answer from the options given below

क

- (1) Both Statement I and Statement II are true
(2) Statement I is true but Statement II is false
(3) Both Statement I and Statement II are false
(4) Statement I is false but Statement II is true

MATRIX JEE ACADEMY**Office : Piprali Road, Sikar (Raj.) | Ph. 01572-241911****Website : www.matrixedu.in ; Email : smd@matrixacademy.co.in**









Ans. Official answer NTA (3)

Sol.

Question ID : 444792667

64. From the following, how many compounds contain at least one secondary alcohol?



Choose the correct answer from the options given below:

क

- (1) Three (2) Four (3) Five (4) Two

Ans. Official answer NTA (1)

Sol.

Question ID : 444792651

65. One mole of $\text{Cl}_2(\text{g})$ was passed into 2 L of cold 2 M KOH solution. After the reaction, the concentrations of Cl^- , ClO^- and OH^- are respectively (assume volume remains constant)

क

- (1) 0.75M, 0.75M, 1M
(2) 0.5M, 0.5M, 1M
(3) 1M, 1M, 1M
(4) 0.5M, 0.5M, 0.5M

Ans. Official answer NTA (2)

Sol.

Question ID : 444792656

66. Two liquids A and B form an ideal solution at temperature T K. At T K, the vapour pressures of pure A and B are 55 and 15kNm^{-2} respectively. What is the mole fraction of A in solution of A and B in equilibrium with a vapour in which the mole fraction of A is 0.8 ?

क

- (1) 0.5217 (2) 0.340 (3) 0.663 (4) 0.480

Ans. Official answer NTA (1)

Sol.

Question ID : 444792653

67. Pair of species among the following having same bond order as well as paramagnetic character will be-

क

- (1) O_2^- , N_2^-
(2) O_2^+ , N_2^-
(3) O_2^- , N_2^+
(4) O_2^+ , N_2^{2-}

Ans. Official answer NTA (2)

Sol.

Question ID : 444792662

68. Find out the statements which are not true.

- A. Resonating structures with more number of covalent bonds and lesser charge separation are more stable.
- B. In electromeric effect, an unsaturated system shows +E effect with nucleophile and -E effect with electrophile.
- C. Inductive effect is responsible for high melting point, boiling point and dipole moment of polar compounds.
- D. The greater the number of alkyl groups attached to the doubly bonded carbon atoms, higher is the heat of hydrogenation.
- E. Stability of carbanion increases with the increase in s – character of the carbon carrying the negative charge.

Choose the correct answer from the options given below:

क

- (1) A, D & E only (2) B & D only

MATRIX JEE ACADEMY**Office : Piprali Road, Sikar (Raj.) | Ph. 01572-241911****Website : www.matrixedu.in ; Email : smd@matrixacademy.co.in**

(3) A, C & D only

(4) B, D & E only

Ans. Official answer NTA (2)

Sol.

Question ID : 444792666

69. A student has planned to prepare acetanilide from aniline using acetic anhydride.

The student has started from 9.3 g of aniline. However, the student has managed to obtain 11 g of dry acetanilide.

The % yield of this reaction is :-

क

(1) 59.5%

(2) 81.5%

(3) 97.5%

(4) 72.5%


Ans. Official answer NTA (2)

Sol.


Question ID : 444792663

70. Given below are two statements:

Statement I: There are several conformers for n-butane. Out of those conformers,



is the least stable and most stable conformer is



Statement II: As the dihedral angle increases, torsional strain decreases from (X) to (Y).

In the light of the above statements, choose the correct answer from the options given below

क

(1) Both Statement I and Statement II are false

(2) Statement I is false but Statement II is true

(3) Both Statement I and Statement II are true

(4) Statement I is true but Statement II is false

Ans. Official answer NTA (3)

Sol.
MATRIX JEE ACADEMY

Office : Piprali Road, Sikar (Raj.) | Ph. 01572-241911

Website : www.matrixedu.in ; Email : smd@matrixacademy.co.in

SECTION - B

Question ID : 444792672

71. Grignard reagent RMgBr(P) reacts with water and forms a gas (Q). One gram of Q occupies 1.4dm^3 at STP. (P) on reaction with dry ice in dry ether followed by H_3O^+ forms a compound (Z). 0.1 mole of (Z) will weigh _____ g. (Nearest integer)

क

Ans. Official answer NTA (6)**Sol.**

Question ID : 444792675

72. The half-life of ^{65}Zn is 245 days. After x days, 75% of original activity remained.

The value of x in days is _____ (Nearest integer)

(Given: $\log 3=0.4771$ and $\log 2=0.3010$)

क

Ans. Official answer NTA (102)**Sol.**

Question ID : 444792671

73. A chromium complex with a formula $\text{CrCl}_3 \cdot 6\text{H}_2\text{O}$ has a spin only magnetic moment value of 3.87 BM and its solution conductivity corresponds to 1 : 2 electrolyte. 2.75 g of the complex solution was initially passed through a cation exchanger. The solution obtained after the process was reacted with excess of AgNO_3 . The amount of AgCl formed in the above process is _____ g. (Nearest integer)

[Given: Molar mass in gmol^{-1} Cr : 52; Cl : 35.5, Ag : 108, O : 16, H : 1]

क

Ans. Official answer NTA (3)**Sol.**

Question ID : 444792673

74. 0.25 g of an organic compound "A" containing carbon, hydrogen and oxygen was analysed using the combustion method. There was an increase in mass of CaCl_2 tube and potash tube at the end of the experiment. The amount was found to be 0.15 g and 0.1837 g, respectively. The percentage of oxygen in compound A is _____ %. (Nearest integer) (Given: molar mass in gmol^{-1} H : 1, C : 12, O : 16)

क

Ans. Official answer NTA (73)**Sol.**

Question ID : 444792674

75. Molar conductivity of a weak acid HQ of concentration 0.18 M was found to be $1/30$ of the molar conductivity of another weak acid HZ with concentration of 0.02 M. If $\lambda^\circ_{Q^-}$ happened to be equal with $\lambda^\circ_{Z^-}$, then the difference of the pK_a values of the two weak acids ($pK_a(\text{HQ}) - pK_a(\text{HZ})$) is _____ (Nearest integer).

[Given: degree of dissociation (α) $\ll 1$ for both weak acids, λ° : limiting molar conductivity of ions]

क

Ans. Official answer NTA (2)**Sol.**